

A DISTRIBUTION OF THE TEMPERATURE IN A RING OF THE INCOMPRESSIBLE, VISCOUS LIQUID WITH TWO FREE BOUNDARIES. EXACT SOLUTIONS.

Belmetsev N.F., Bytev V.O.
Tyumen State University, Russia

e-mail: weqsmachine@gmail.com, vbytev@utmn.ru

October, 2009

Abstract. It has been found the exact solutions for nonstationary distribution of the temperature in the liquid ring with two viscosities (μ and μ_0) and two free boundaries of the ring.

We consider the problem about rotational symmetric movement of viscous and nonviscous liquid in a ring for nonclassical model of hydrodynamics (see [1, 2]). A similar problem for classical models Navier-Stokes and Stokes was considered in papers [3, 4, 5, 6, 7]. Previous papers contain the proves for theorems of global resolvability and uniqueness, and contain the asymptotes for solutions behavior. In the moment $t = 0$, we see that the liquid fills a ring $R_{20} < r < R_{10}$ and has distribution of velocities:

$$u_r = \varphi_0 r^{-1}, \quad u_\theta = v_0(r).$$

Here, $R_{10}, R_{20} > 0$, and φ_0 are constants, $v_0(r)$ is an arbitrary smooth function. Since u_θ is not depend from θ , we see that the specified association u_r from r is unique.

Let R_{10}, R_{20} are free boundaries. In this case the liquid ring begins movement by inertia. Since we have rotational symmetry of an initial condition, we may assume that this one is similarly in all time agreement. Using this statement, we get that free boundaries are remained circles during all time of movement: $r = R_1(t)$, $r = R_2(t)$.

Let us note the equations of movement and thermal conduction for nonclassical liquid [1, 2] with the supposition of its incompressibility and homogeneity:

$$\begin{aligned} \frac{\partial \vec{u}}{\partial t} + (\vec{u} \cdot \nabla) \vec{u} - \frac{1}{\rho} M \Delta \vec{u} + \frac{1}{\rho} \nabla p &= 0 \\ \operatorname{div} \vec{u} &= 0 \\ \rho C_p \frac{dT}{dt} &= k \Delta T + \rho \nu H \end{aligned}$$

Here, $\vec{u}$ is a vector of velocity with components u_i ; p is a hydrostatic (equilibrium) pressure; ρ is a density; H is a dissipative function; C_p is a thermal capacity; k is a factor of thermal conduction; $\mu = \rho \nu$ is a viscosity; M is a tensor of viscosity:

$$M = \begin{pmatrix} \mu & \mu_0 \\ -\mu_0 & \mu \end{pmatrix}$$

μ_0 is nondissipative viscosity which can accept values with any signs. The stress tensor:

$$T = -pI + 2MD$$

D is the tensor with components:

$$D_{ij} = \frac{1}{2} \left(\frac{\partial u_i}{\partial x_j} + \frac{\partial u_j}{\partial x_i} \right), \quad (i, j = 1, 2).$$

Let $r = R_1(t)$ be an exterior boundary of the ring and $r = R_2(t)$ be an interior boundary of the ring. Since,

$$u_r = u(r, t); \quad u_\theta = v(r, t); \quad p = p(r, t),$$

then a components of the stress tensor in polar coordinates are:

$$\begin{aligned} T_{rr} &= -p + 2\mu \frac{\partial u}{\partial r} + \mu_0 \left(\frac{\partial v}{\partial r} - \frac{v}{r} \right); \\ T_{r\theta} &= \mu \left(\frac{\partial v}{\partial r} - \frac{v}{r} \right) - 2\mu_0 \frac{\partial u}{\partial r}; \\ T_{\theta\theta} &= -p + 2\mu \frac{u}{r} - \mu_0 \left(\frac{\partial v}{\partial r} - \frac{v}{r} \right). \end{aligned}$$

Using this representation, we obtain:

$$\begin{aligned}
u &= \frac{\Phi(t)}{r}; \\
\frac{1}{r} \frac{d\Phi}{dt} - \frac{\Phi^2}{r^3} - \frac{v^2}{r} &= -\frac{\partial}{\partial r} \left(p - \nu_0 \frac{1}{r} \frac{\partial(rv)}{\partial r} \right); \\
\frac{\partial v}{\partial t} + \frac{\Phi}{r} \frac{\partial v}{\partial r} + \frac{\Phi v}{r^2} &= \nu \left(\frac{\partial^2 v}{\partial r^2} + \frac{1}{r} \frac{\partial v}{\partial r} - \frac{v}{r^2} \right); \\
\rho C_p \left(\frac{\partial T}{\partial t} + u \frac{\partial T}{\partial r} + \frac{v}{r} \frac{\partial T}{\partial \theta} \right) &= k \left[\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial T}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 T}{\partial \theta^2} \right] + \\
&+ \mu \left\{ 2 \left(\frac{\partial u}{\partial r} \right)^2 + 2 \left(\frac{u}{r} \right)^2 + \left[r \frac{\partial}{\partial r} \left(\frac{v}{r} \right) \right]^2 \right\}.
\end{aligned}$$

Here u is the radial component of velocity; v is the azimuthal component of velocity. Since the stress vector is equal to zero on free boundaries, we get boundary conditions:

$$T_{rr} = - \left(p + \frac{2\nu\Phi(t)}{r^2} \right) + \nu_0 \left(\frac{\partial v}{\partial r} - \frac{v}{r} \right) = 0, \text{ if } r = R_{1,2}(t);$$

$$T_{r\theta} = \nu r \frac{\partial}{\partial r} \left(\frac{v}{r} \right) - 2\nu_0 \frac{\Phi(t)}{r^2} = 0, \text{ if } r = R_{1,2}(t).$$

Also, we have dynamic condition on free boundaries:

$$\frac{dR_i(t)}{dt} = \frac{\Phi(t)}{R_i(t)}, \quad (i = 1, 2)$$

Supposing $\mu_0^2 + \mu^2 \neq 0$, we have:

$$\begin{aligned}
\frac{\partial v}{\partial t} + \frac{\Phi}{r} \frac{\partial v}{\partial r} + \frac{\Phi v}{r^2} &= \nu \frac{\partial}{\partial r} \left[\frac{1}{r} \frac{\partial(rv)}{\partial r} \right]; \\
\nu \left(\frac{\partial v}{\partial r} - \frac{v}{r} \right) + 2\nu_0 \frac{\Phi(t)}{r^2} &= 0, \quad r = R_{1,2}(t) \\
\frac{d\Phi}{dt} \ln \left(\frac{R_1}{R_2} \right) &= \int_{R_2(t)}^{R_1(t)} \left[\frac{v^2}{r} + 2\nu_0 \frac{\partial}{\partial r} \left(\frac{v}{r} \right) \right] dr - \left(2\nu\Phi - \frac{\Phi^2}{2} \right) \left[\frac{1}{R_2^2} - \frac{1}{R_1^2} \right]; \\
\Phi(0) &= \Phi_0; \\
v(r, 0) &= v_0(r);
\end{aligned}$$

The equality

$$R_1^2(t) - R_2^2(t) = R_{10}^2 - R_{20}^2;$$

is the conservation law of square. The equality

$$\int_{R_2(t)}^{R_1(t)} r^2 v(r, t) dr = \int_{R_{20}}^{R_{10}} r^2 v(r, 0) dr \equiv \int_{R_{20}}^{R_{10}} r^2 v_0(r) dr;$$

is the conservation law of impulse moment.

Let us write down the equation for temperature T :

$$\begin{aligned}
\rho C_p \left(\frac{\partial T}{\partial t} + u \frac{\partial T}{\partial r} + \frac{v}{r} \frac{\partial T}{\partial \theta} \right) &= k \left[\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial T}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 T}{\partial \theta^2} \right] + \\
&+ \mu \left\{ 2 \left(\frac{\partial u}{\partial r} \right)^2 + 2 \left(\frac{u}{r} \right)^2 + \left[r \frac{\partial}{\partial r} \left(\frac{v}{r} \right) \right]^2 \right\};
\end{aligned}$$

$$T|_{t=0} = T_0(r, \theta).$$

Boundary conditions for T will be specified additionally.

Let

$$\begin{aligned} \tau &= \frac{\nu}{R_{20}^2} t; & \eta &= \frac{r^2 - R_2^2(t)}{R_{20}^2}; & \xi &= \frac{R_2^2(t)}{R_{20}^2}; \\ u &= \frac{\nu \Psi}{r}; & v &= \frac{\nu}{R_{20}^2} r \omega; & \Psi &= \frac{\Phi}{\nu}; & a &= \frac{R_{10}^2}{R_{20}^2} - 1; \\ \Theta &= \frac{T}{T_0}; & \Theta_0 &= T_0 \end{aligned}$$

be a new coordinates then we have a problem:

$$\frac{\partial \omega}{\partial \tau} + \frac{2\Psi}{\xi + \eta} \omega = 4(\xi + \eta) \frac{\partial^2 \omega}{\partial \eta^2} + 8 \frac{\partial \omega}{\partial \eta}; \quad (1)$$

$$\omega|_{\tau=0} = \omega_0(\eta), \quad 0 \leq \eta \leq a; \quad (2)$$

$$\frac{\partial \omega}{\partial \eta} + \varepsilon \frac{\Psi}{(\xi + \eta)^2} = 0, \quad \text{if } \eta = 0, a; \quad \tau \geq 0; \quad (3)$$

$$\frac{d\omega}{d\tau} = \frac{a\Psi(\Psi - 4)}{\xi(\xi + a) \ln(1 + \frac{a}{\xi})} + \frac{1}{\ln(1 + \frac{a}{\xi})} \int_0^a \left(\omega^2 + 4\varepsilon \frac{\partial \omega}{\partial \eta} \right) d\eta; \quad (4)$$

$$\frac{d\xi}{d\tau} = 2\Psi; \quad \Psi(0) = \Psi_0; \quad \xi(0) = 1; \quad \varepsilon = \frac{\nu_0}{\nu}; \quad (5)$$

$$\begin{aligned} & \frac{\rho C_p R_{20}^2 \Theta_0}{4\mu^2} \left(\frac{\partial \Theta}{\partial \tau} + \omega \frac{\partial \Theta}{\partial \theta} \right) = \\ &= \frac{k R_{20}^2 \Theta_0}{4\mu^3} \left(\frac{\partial \Theta}{\partial \eta} + (\xi + \eta) \frac{\partial^2 \Theta}{\partial \eta^2} + \frac{1}{4(\xi + \eta)} \frac{\partial^2 \Theta}{\partial \theta^2} \right) + \\ &+ \frac{\Psi^2}{(\xi + \eta)^2} + (\xi + \eta)^2 \left(\frac{\partial \omega}{\partial \eta} \right)^2 \end{aligned} \quad (6)$$

It is easy to see that

$$\omega = \frac{4\varepsilon}{\xi + \eta}; \quad \omega_0(\eta) = \frac{4\varepsilon}{1 + \eta}; \quad \Psi = \Psi_0 = 4; \quad \xi(\tau) = 8\tau + 1 \quad (7)$$

is an exact solution of the problem (1)–(5). Using (6),(7), and $\Theta = \Theta(\tau(t), \eta(r, t))$, we get:

$$A \frac{\partial \Theta}{\partial \tau} = B \frac{\partial}{\partial \eta} \left((8\tau + \eta + 1) \frac{\partial \Theta}{\partial \eta} \right) + \frac{16(1 + \varepsilon^2)}{(8\tau + \eta + 1)^2} \quad (8)$$

Here

$$A = \frac{\rho C_p R_{20}^2 \Theta_0}{4\mu^2}, \quad B = \frac{k R_{20}^2 \Theta_0}{4\mu^3}, \quad \varepsilon = \frac{\nu_0}{\nu} \quad (9)$$

are dimensionless parameters.

Using methods of Group Analysis (see [9, 10, 11, 12]), we obtain infinitesimal operator:

$$X = \xi^1 \frac{\partial}{\partial \tau} + \xi^2 \frac{\partial}{\partial \eta} + \eta^1 \frac{\partial}{\partial \Theta} \quad (10)$$

with coefficients:

$$\begin{cases} \xi^1 = C_1 \tau^2 / 2 + C_2 \tau + C_3; \\ \xi^2 = C_1 \tau (4\tau + \eta + 1) + C_2 (\eta + 1) - 8 C_3; \\ \eta^1 = (C_4 - C_1 (\tau + (A/B)\eta) / 2) \Theta + b_2(\tau, \eta); \end{cases} \quad (11)$$

Here C_i are arbitrary real constants ($i = 1, 2, 3, 4$). A function $b_2(\tau, \eta)$ should be a solution of following equation:

$$\begin{aligned} A \frac{\partial b_2}{\partial \tau} = B \left(\frac{\partial b_2}{\partial \eta} + (8\tau + \eta + 1) \frac{\partial^2 b_2}{\partial \eta^2} \right) + \\ + \frac{16(1 + \varepsilon^2)}{(8\tau + \eta + 1)^2} \left(-(C_2 + C_4) - \frac{C_1}{2} \left(\tau - \frac{A}{B} \eta \right) \right) \end{aligned} \quad (12)$$

Supposing that $C_1 = 0$, $C_2 = 0$, $C_4 = 0$, and $b_2(\tau, \eta) = 0$, we get:

$$\begin{aligned} X = C_3 \frac{\partial}{\partial \tau} - 8C_3 \frac{\partial}{\partial \eta} \\ \begin{cases} I_1 = 8\tau + \eta + 1 \\ I_2 = \Theta \end{cases} \end{aligned}$$

Using the theorem for representation of invariant solutions through invariants from [9], we receive for unknown function Θ :

$$\Theta = I_2 = \varphi(I_1) = \varphi(8\tau + \eta + 1)$$

Combining this with (8) we get the ordinary differential equation:

$$B I_1 \frac{d^2 \varphi(I_1)}{d I_1^2} + (B - 8A) \frac{d \varphi(I_1)}{d I_1} + \frac{16(1 + \varepsilon^2)}{(I_1)^2} = 0$$

Solving this equation, and checking solution by input equation we obtain the invariant solution of equation (8):

$$\Theta(\tau, \eta) = C_5 - \frac{16(1 + \varepsilon^2)}{(8\tau + \eta + 1)(B + 8A)} \quad (13)$$

Let $C_1 = 0$, $C_2 = 1$, $C_4 = -2$, and function $b_2(\tau, \eta)$ is the partial solution (13):

$$b_2(\tau, \eta) = C_5 - \frac{16(1 + \varepsilon^2)}{(8\tau + \eta + 1)(B + 8A)}$$

then for infinitesimal operator (10) (and for invariants), we obtain:

$$\begin{aligned} X = (\tau + C_3) \frac{\partial}{\partial \tau} + (1 + \eta - 8C_3) \frac{\partial}{\partial \eta} + (b_2(\tau, \eta) - 2\Theta) \frac{\partial}{\partial \Theta} \\ \begin{cases} J_1 = (1 + \eta - 8C_3)/(\tau + C_3) \\ J_2 = (\tau + C_3)((\tau + C_3)\Theta - \tau C_5/2) - \tau C_3 C_5/2 + \\ + 16\tau(\tau + C_3)(1 + \varepsilon^2)/((8\tau + \eta + 1)(8A + B)) \end{cases} \end{aligned}$$

Using J_1 , J_2 , we have (in system of computer calculus Maple) the invariant solution for equation (8):

$$\begin{aligned} \Theta(\tau, \eta) = K \left(\frac{A(1 + \eta - 8C_3) - B(\tau + C_3)}{(\tau + C_3)^3} \right) \times \\ \times \exp \left\{ -\frac{A(1 + \eta - 8C_3)}{B(\tau + C_3)} \right\} \left(\frac{8\tau + \eta + 1}{\tau + C_3} \right)^{\frac{8A}{B}} + \\ + \frac{C_5}{2} - \frac{16(1 + \varepsilon^2)}{(8\tau + \eta + 1)(B + 8A)}, \quad (14) \\ K = \frac{8A K_1 + B(K_1 - K_2)}{B(8A + B)} = const. \end{aligned}$$

Here K_1 , K_2 are constants. It is easy to see that the expressions (13), (14) are solutions for equation (8), and:

$$\lim_{\tau \rightarrow +\infty} (\Theta(\tau, \eta)) = \frac{C_5}{2}.$$

Let

$$\begin{cases} \Theta_1(\tau) = \Theta(\tau, a) \\ \Theta_2(\tau) = \Theta(\tau, 0) \end{cases} \quad (15)$$

be a boundary conditions.

Using (14) and (15) we receive:

$$\begin{cases} \Theta_1(\tau) = K \left(\frac{A(1+a-8C_3)-B(\tau+C_3)}{(\tau+C_3)^3} \right) \exp \left\{ -\frac{A(1+a-8C_3)}{B(\tau+C_3)} \right\} \left(\frac{8\tau+a+1}{\tau+C_3} \right)^{\frac{8A}{B}} + \\ + \frac{C_5}{2} - \frac{16(1+\varepsilon^2)}{(8\tau+a+1)(B+8A)} \\ \Theta_2(\tau) = K \left(\frac{A(1-8C_3)-B(\tau+C_3)}{(\tau+C_3)^3} \right) \exp \left\{ -\frac{A(1-8C_3)}{B(\tau+C_3)} \right\} \left(\frac{8\tau+1}{\tau+C_3} \right)^{\frac{8A}{B}} + \\ + \frac{C_5}{2} - \frac{16(1+\varepsilon^2)}{(8\tau+1)(B+8A)} \end{cases} \quad (16)$$

The difference of temperatures on boundaries at $\tau = 0$ is defined by the constant:

$$\begin{aligned} C = \Theta_1(0) - \Theta_2(0) &= \frac{16a(1+\varepsilon^2)}{(1+a)(8A+B)} + \frac{K}{C_3^3} \exp \left\{ -\frac{A(1-8C_3)}{B C_3} \right\} \times \\ &\times \left((A(1+a-8C_3) - B C_3) \exp \left\{ -\frac{Aa}{B C_3} \right\} \left(\frac{1+a}{C_3} \right)^{\frac{8A}{B}} - (A(1-8C_3) - B C_3) \left(\frac{1}{C_3} \right)^{\frac{8A}{B}} \right) \end{aligned}$$

We have distribution of temperature $\Theta(\tau, \eta)$ for $\tau = 0$:

$$\begin{aligned} \Theta(0, \eta) &= K \left(\frac{A(1+\eta-8C_3)-B C_3}{C_3^3} \right) \exp \left\{ -\frac{A(1+\eta-8C_3)}{B C_3} \right\} \left(\frac{\eta+1}{C_3} \right)^{\frac{8A}{B}} + \\ &+ \frac{C_5}{2} - \frac{16(1+\varepsilon^2)}{(\eta+1)(B+8A)}, \end{aligned} \quad (17)$$

Let us choose constants as follows:

$$\begin{aligned} C = 0, \quad A = \frac{3}{4}, \quad B = 6, \quad \varepsilon = \frac{1}{2}, \quad a = 1, \quad C_3 = \frac{1}{8}, \\ K = \frac{\frac{10}{9}C_3^4}{4(8C_3 - 1) \exp \left\{ 1 - \frac{1}{4C_3} \right\} - (16C_3 - 1) \exp \left\{ 1 - \frac{1}{8C_3} \right\}} = -\frac{5}{6^2 8^3} \approx -0.00027127 \end{aligned} \quad (18)$$

Values of temperature on boundaries at $\tau = 0$ with such choice of constants are identical.

Considering values of constants (18) and expressions for temperature on boundaries (16), we have the following problem:

$$\frac{\partial \Theta}{\partial \tau} - \frac{\partial}{\partial \eta} \left(8(8\tau + \eta + 1) \frac{\partial \Theta}{\partial \eta} \right) - \frac{80}{3(8\tau + \eta + 1)^2} = 0 \quad (19)$$

$$\begin{cases} \left. \frac{\partial \Theta}{\partial \eta} \right|_{\eta=0} = \frac{5(1+16\tau)}{6(8\tau+1)^3} \\ \left. \frac{\partial \Theta}{\partial \eta} \right|_{\eta=a} = \left(\frac{5}{3(8\tau+2)^2} \right) + \left(\frac{5}{6(8\tau+2)^3} \right) \left(\frac{1-(8\tau+1)(8\tau+3)}{(8\tau+1)^2} \right) \end{cases} \quad (20)$$

$$\Theta(0, \eta) = \frac{C_5}{2} - \left(\frac{5}{6} \right) \left((\eta^2 - 1) \exp(-\eta) + \frac{2}{(\eta + 1)} \right) \quad (21)$$

Substituting values of constants (18) in expression (14), we receive the solution of the problem (19)-(21):

$$\Theta(\tau, \eta) = \frac{C_5}{2} - \left(\frac{5}{6} \right) \left(\frac{2}{(8\tau + \eta + 1)} + \left(\frac{\eta^2 - (8\tau + 1)^2}{(8\tau + 1)^4} \right) \exp \left\{ -\frac{\eta}{8\tau + 1} \right\} \right) \quad (22)$$

Supposing temperature nonnegative, we get restriction on value of a free constant: $C_5 \geq 5/3$.

Considering (7), we get the solution (14) in (t, r) co-ordinates:

$$T(t, r) = T_0 K \left(\frac{AR_{20}^4 r^2 - (8A + B)(\nu R_{20}^4 t + C_3 R_{20}^6)}{(\nu t + C_3 R_{20}^2)^3} \right) \times$$

$$\times \exp \left\{ \frac{8A}{B} - \frac{Ar^2}{B(\nu t + C_3 R_{20}^2)} \right\} \left(\frac{r^2}{(\nu t + C_3 R_{20}^2)} \right)^{\frac{8A}{B}} + \\ + \frac{C_5 T_0}{2} - \frac{16 T_0 R_{20}^2 (1 + \varepsilon^2)}{r^2 (B + 8A)}.$$

Also, we have:

$$R_{20}^2(t) = 8\nu t + R_{20}^2, \quad R_{10}^2(t) = 8\nu t + R_{10}^2, \\ u = \frac{4\nu}{r}, \quad v = \frac{4\nu_0}{r}.$$

As a result, exact solutions are received, the general view of Lie group for the equation (19) is received, the asymptote is found, and profiles for one of the solutions are constructed. These results allows to build another solutions for a problem about temperature distribution in the liquid ring.

References

- [1]. *Andreev V.K., Bublik V.V., Bytev V.O.* Symmetry of nonclassical hydrodynamics models. Novosibirsk, Nauka, 2003 (in Russian).
- [2]. *Bytev V.O.* The simple nonpolar continuum media. Part II. The constitutive eqations. Linear structures. // (archived article, <http://www.arxiv.org>), 2009.
- [3]. *Ovsyannikov L.V.* Equations and examples. Novosibirsk, Nauka, 1967 (in Russian).
- [4]. *Bytev V.O.* Nonstationary moving of viscous incompressible liquid ring with free boundary // PMTF 3, 1970, p. 88–98 (in Russian).
- [5]. *Pukhnachev V.V.* Nonclassical problem of a boundary layer theory. Novosibirsk, NGU, 1980 (in Russian).
- [6]. *Lavrenti'eva O.M.* Limiting modes of movement of a rotating viscous ring. DSS, Novosibirsk, 1980. No. 4. p. 15–34.
- [7]. *Lavrenti'eva O.M.* Nonstationary moving of rotating viscous capillary liquid's ring. DSS, Novosibirsk, 1978, No. 31. p. 52–60 (in Russian).
- [8]. *Sedov L.I.* Similarity and Dimensional Methods in Mechanics. Moscow, Nauka, 1977 (in Russian).
- [9]. *Ovsyannikov L.V.* Group properties of differential equations. Academic Press, 1962 (in Russian).
- [10]. *Ibragimov N.H.* An ABC of group analysis. Novoe v Zhizni, Nauke, Tekhnike. Seriya Matematika, Kibernetika, No. 8, Znanie, Moscow, 1989 (in Russian).
- [11]. *Ibragimov N.H.* A Practical Course in Differential Equations and Mathematical Modelling. Second edition. ALGA Publications, Blekinge Institute of Technology Karlskrona, Sweden, 2005.
- [12]. *Meleshko, S.V.* Methods for constructing exact solutions of partial differential equations: mathematical and analytical techniques with applications to engineering / S.V. Meleshko. p. cm. - (Mathematical and analytical techniques with applications to engineering), - New York, USA: Springer Science + Business Media, Inc. 2005.